

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



JBP 438

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 38/08, 38/48, 38/55, 38/17, 38/05 // 7/00, 9/127</b>	<b>A1</b>	(11) International Publication Number: <b>WO 99/30729</b> (43) International Publication Date: <b>24 June 1999 (24.06.99)</b>
---	-----------	--

<p>(21) International Application Number: <b>PCT/US98/26255</b></p> <p>(22) International Filing Date: <b>10 December 1998 (10.12.98)</b></p> <p>(30) Priority Data: <b>60/069,797</b>                      <b>16 December 1997 (16.12.97)</b>      <b>US</b></p> <p>(71) Applicant: <b>JOHNSON &amp; JOHNSON CONSUMER COMPANIES, INC. [US/US]; Grandview Road, Skillman, NJ 08558 (US).</b></p> <p>(72) Inventors: <b>MIRI, Sciberg, 168 Herrontown Road, Princeton, NJ 08540 (US). SHAPIRO, Stanley, S.; 10 Plymouth Drive, Livingston, NJ 07039 (US). EISINGER, Magdalena; 30 Pine Terrace, Demarest, NJ 07627 (US).</b></p> <p>(74) Agents: <b>CIAMPORCERO, Audley, A. et al.; Johnson and Johnson, One Johnson &amp; Johnson Plaza, New Brunswick, NJ 08933 (US).</b></p>	<p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b></p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
--	---

(54) Title: **COMPOSITIONS AND METHODS FOR REGULATING PHAGOCYTOSIS AND ICAM-1 EXPRESSION**

(57) Abstract

This invention provides compositions of matter for treating and preventing certain mammalian disorders ameliorated by either an increase or decrease in phagocytosis or ICAM-1 expression in appropriate cells. This invention also provides method of altering the phagocytosis or ICAM-1 expression level in a cell. This invention further provides methods of treating and preventing mammalian disorders affected by the alteration of phagocytosis or ICAM-1 expression. The instant methods and composition of matter all relate to the use of agents that specifically increase or decrease phagocytosis or ICAM-1 expression. Finally, this invention provides related articles of manufacture.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5 COMPOSITIONS AND METHODS FOR REGULATING PHAGOCYTOSIS AND  
ICAM-1 EXPRESSION

Field of the Invention

10 This invention relates to the prevention and  
treatment of mammalian disorders that are ameliorated  
by altering phagocytosis or ICAM-1 expression in  
certain cells. The invention provides numerous  
compositions, methods and articles of manufacture, and  
addresses a considerable range of disorders such as  
15 those of skin and the immune and central nervous  
systems. This invention is based on the discovery of a  
mechanism for the regulation of phagocytosis and ICAM-1  
expression.

20 Background of the Invention

Phagocytosis and ICAM-1 Expression Generally

25 Phagocytosis is the cellular process of ingestion,  
and usually of isolation or destruction, of particulate  
material. In vertebrates, it is a characteristic  
function of various leukocytes and reticuloendothelial  
cells. Phagocytosis serves as an important bodily  
defense mechanism against infection by microorganisms,  
30 and against occlusion of mucous surfaces and tissues by  
foreign particles and tissue debris. Phagocytosis is  
distinct from pinocytosis, which is the uptake of fluid  
by a cell through invagination and pinching off of the  
plasma membrane. Herein, the terms "phagocytosis" and  
35 "cellular ingestion" are used interchangeably.

Intercellular adhesion molecule -1 ("ICAM-1") is an inducible cell-surface glycoprotein that is implicated in cell-cell adhesion and phagocytosis. In particular, the regulation of ICAM-1 plays a role in inflammatory situations, septic shock and neurological disorders (reviewed in van de Stolpe and van der Saag, J Mol Med 74:1, 13-33, 1996). ICAM-1 is elevated in autoimmune diseases such as rheumatoid arthritis and psoriasis. Inflammatory and immune responses are impaired in mice deficient in ICAM-1 (Sligh et al., PNAS 90:8529-33, 1993).

#### Mammalian Disorders Related to Phagocytosis and ICAM-1 Expression

The levels of phagocytosis and ICAM-1 expression in different cells have important implications. Numerous examples of these implications are provided here.

#### *Immune-Related and Inflammatory Disorders*

The primary cause of pulmonary emphysema is the accumulation of foreign material (e.g. smoke condensate) in the lung. This accumulation is followed by the recruitment of neutrophils that are degranulated during attempted phagocytosis (Travis, et. al., Am. J. Respir. Crit. Care Med. Vol. 150:5143-5146, 1994).

Immunological lung disorders such as allergic bronchopulmonary aspergillosis cause mucus plugging of airways, eosinophylic pneumonia and bronchiolitis obliterans. In such diseases, neutrophil elastase-cleaved immunoglobulins and digested C3b receptors limit the phagocytosis of pathogens (Greenberger, JAMA, Vol. 278, No.22, 1997). The increase in neutrophil elastase,

whil impairing phagocyt sis, is beneficial for fighting persistent bacterial infections in the lungs, especially in CF patients (Doring, et al., Am. J. Respir. Crit. Care Med. 150:6 Pt 2, S114-7, 1994).

5

Periodontal diseases start with the accumulation of plaque at the base of the teeth, followed by the growth of opportunistic bacteria below the gum line. As with the immune response in emphysema, neutrophils are recruited to the infected site, followed by their degranulation during frustrated phagocytosis (Travis, et al., Am. J. Respir. Crit. Care Med. Vol. 150:5143-5146, 1994). The rates of adhesion and ingestion of opsonized Staphylococcus Aureus by polymorphonuclear cells ("PMN's") from periodontal patients is significantly reduced relative to healthy controls (MacFarlane, et al., J Periodontol 1992; 63:908-913, 1992).

20 Individuals who are genetically immuno-compromised, who have acquired immuno-suppression (such as HIV-infected individuals), or who have temporarily acquired immuno-suppression (such as that following organ transplantation, foreign implants, valve replacement or 25 cancer treatment, and the like), often suffer from secondary infections.

Pulmonary polymorphnuclear leukocytes from diabetic patients were shown to have reduced phagocytic 30 activities, both at the level of ingestion and killing of bacteria, compared to healthy individuals (e.g. Musclow, et al, Cytobios, 65:15-24 1991). In particular, diabetic abnormalities in the immune response include impaired chemotaxis, impaired phagocytosis and impaired adhesion 35 (Grant-Theule, Periodontal Abstracts, Vol. 44, No. 3,

1996). These patients often suffer from undesired infections.

#### *Cardiovascular System Disorders*

5

The formation of atherosclerotic plaques is induced by aging or by restenosis following balloon angioplasty. Atherosclerotic lesions contain cholesterol-rich particles, many of which aggregate and are internalized in an unregulated fashion by macrophage phagocytosis. This phagocytic process is independent of the LDL or scavenger receptor. The lipid-loaded macrophages, called foamy cells, can lead to further growth of the atherosclerotic plaque (Hoff, et al., *European Heart Journal*, II (Supp. E), 105-115, 1990; Robert, et al., *Annals New York Acad. of Sciences*, 673:331-341, 1992).

#### *Central Nervous System Disorders*

20 Microglial cells found at the periphery of amyloid plaque cores have been shown to contain plaque fibrils of beta/A4 amyloid (El Hachimi and Foncin, *C.R. Acad. Sci. Paris, Sciences de la vie/Life sciences*, 317:445-451, 1994). The ability of microglial cells to phagocytose and clear senile plaque cores is suppressed in the presence of an astrocyte-secreted diffusable factor. This factor prevents the clearance of senile plaques, allowing them to persist in Alzheimer's disease and other neuropathological degenerative processes (DeWitt, et al., *Experimental Neurology*, 149:329-340, 1998).

Neutrophil phagocytosis was found to be reduced in mentally depressed patients (e.g. McAdams and Leonard, *Prog. Neuro-Psychopharmacol. & Biol. Psychiat.*, Vol. 17:971-984, 1993; Maes et al., *J. Psychiat. Res.*, Vol. 35

26, No. 2, 125-134, 1992). Patients with phobic disorders have reduced phagocytosis and cell-killing capacities. Benzodiazepine compounds, used in the treatment of neurological disorders, were shown to reduce or inhibit  
5 phagocytosis (e.g. Covelli et al., *Immunopharmacology and Immunotoxicology*, 11(4):701-714, 1989).

#### *Skin Disorders*

10 Mid-dermal elastosis, a skin disorder, is clinically characterized by the appearance of wrinkles and aged appearance which results, in part, from phagocytosis of morphologically normal elastic tissue (e.g. Fimiani, et al., *Arch Dermatol Res.*, 287:152-157, 1995).

15

Many types of pigmentation disorders exist in diverse forms. These can be inherited (e.g. vitiligo), acquired (e.g. post-inflammatory pityriasis alba, idiopathic guttate hypomelanosis, melasma), and  
20 transmitted through infection (e.g. tinea versicolor). These disorders can be benign and self-limiting (e.g. isolated café au lait spots, photocontact dermatitis), or a sign of a more serious underlying disease (e.g. multiple café au lait spots, malignant acanthosis  
25 nigricans) (Hacker, *Postgrad Med* 99:177-86, 1996).

UV irradiation is known to induce an inflammatory condition and an abnormal regulation of ICAM-1 expression. This induction has been documented in the  
30 form of sunburns and side effects of PUVA therapy. PUVA therapy is used for numerous skin disorders such as psoriasis, a disease associated with upregulation of ICAM-1 expression (e.g. Tronnier, et al., *J. Cutan Pathol* 1997, 24:278-85; Ahrns, et al., *PNAS* 1997, 94:6837-41).

35



Acne vulgaris is a multi-stage disorder. The basic acne lesion is the comedo. The second, inflammatory stage when neutrophils are recruited to the comedo area is the reason the disease progresses. Nearly all problems associated with acne result from this inflammatory phase. Neutrophils from tetracycline-treated patients demonstrate slower migration rates toward chemotactic factors and depressed random migration in vitro (e.g. Webster, J. Am. Acad. Dermatol. 1995, 33:247-53).

10

### Protease-Activated Receptors

The Protease-activated receptor-2 ("PAR-2") is a seven transmembrane G-protein-coupled receptor that is related to, but distinct from, the thrombin receptors ("TR's", also named PAR-1, and PAR-3) and PAR-4 in its sequence. Protease-activated receptors are activated proteolytically by an arginine-serine cleavage at the extracellular domain. The newly created N-termini then activate these receptors as tethered ligands. Both receptors can be activated by trypsin, but only the TR's and PAR-4 are activated by thrombin. Only PAR-2 is activated by mast cell tryptase. These receptors can also be activated by the peptides that correspond to their new N-termini, independent of receptor cleavage. SLIGRL, the mouse PAR-2-activating peptide, is equipotent in the activation of the human receptor, as is SLIGKVD, the human activating peptide. (For a review, see Coughlin, PNAS 91:9200-202, 1994; Brass and Molino, Thrombosis and Haemostasis 78:234-41, 1997; Morley, et al., Can. J. Physiol Pharmacol 25:832-41, 1997.) While the function of TR is well documented, the biology of PAR-2 has not yet been fully identified. A role for PAR-2 activation in the inhibition of keratinocyte growth and

differentiation has been recently described by Derian et al., Cell Growth & Differentiation 8:743-749, 1997.

### Summary of the Invention

This invention provides compositions of matter for  
5 treating and preventing certain mammalian disorders.  
These compositions, and related methods, are based on  
the discovery of a mechanism for the regulation of  
phagocytosis and ICAM-1 expression. The instant  
compositions include the following:

10

(1) a composition of matter for treating a mammal  
afflicted with a disorder ameliorated by an increase in  
phagocytosis or ICAM-1 expression in appropriate cells,  
which comprises (a) a therapeutically effective amount  
15 of an agent that increases phagocytosis or ICAM-1  
expression, and (b) a pharmaceutically or cosmetically  
acceptable carrier;

(2) a composition of matter for treating a mammal  
20 afflicted with a disorder ameliorated by a decrease in  
phagocytosis or ICAM-1 expression in appropriate cells,  
which comprises (a) a therapeutically effective amount  
of an agent that specifically decreases phagocytosis or  
ICAM-1 expression, and (b) a pharmaceutically or  
25 cosmetically acceptable carrier;

(3) a composition of matter for preventing in a  
mammal a disorder ameliorated by an increase in  
phagocytosis or ICAM-1 expression in appropriate cells,  
30 which comprises (a) a prophylactically effective amount  
of an agent that specifically increases phagocytosis or  
ICAM-1 expression, and (b) a pharmaceutically or  
cosmetically acceptable carrier; and

(4) a composition of matter for preventing in a mammal a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a prophylactically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier.

This invention also provides methods of altering the phagocytosis or ICAM-1 expression level in a cell. The invention first provides a method of increasing phagocytosis or ICAM-1 expression in a mammalian cell, comprising contacting the cell with an effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression. Second, this invention provides a method of decreasing phagocytosis or ICAM-1 expression in a mammalian cell, comprising contacting the cell with an effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.

This invention further provides methods of treatment and prophylaxis regarding disorders affected by the alteration of phagocytosis or ICAM-1 expression. Specifically, this invention provides the following:

(1) a method of treating a mammal afflicted with a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a therapeutically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression;

(2) a method of treating a mammal afflicted with a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises

administering to th mammal a therapeutically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression;

5 (3) a method of preventing in a mammal a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a prophylactically effective amount of an agent that specifically  
10 increases phagocytosis or ICAM-1 expression; and

(4) a method of preventing in a mammal a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises  
15 administering to the mammal a prophylactically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.

This invention still further provides an article  
20 of manufacture for administering to a mammal the instant composition of matter, comprising a solid delivery vehicle having the composition operably affixed thereto.

25 Finally, this invention provides a method of administering a therapeutic, prophylactic or cosmetic compound to a mammal, comprising administering to the mammal (a) the compound and (b) a composition of matter comprising a pharmaceutical or cosmetic carrier and an  
30 agent that specifically increases phagocytosis in an amount sufficient to increase phagocytosis in cells where uptake of the compound is desired, wherein the composition is administ red prior to and/or concurrently with the administration of the compound.

**Brief Description of the Figures**

5 Figure 1 shows primary keratinocytes exposed to fluorescent microspheres following treatment with Compound I or SLIGRL.

10 Figure 2 shows cells of a keratinocyte cell line exposed to fluorescent microspheres following treatment with Compound I or SLIGRL.

15 Figure 3 shows cells of a fibroblast cell line exposed to fluorescent microspheres following treatment with soybean trypsin inhibitor ("STI") or SLIGRL.

Figure 4A shows a dose-response graph of macrophages treated with STI and exposed to fluorescent *E. Coli*.

20 Figure 4B shows a dose-response graph of macrophages treated with Compound I or SLIGRL and exposed to fluorescent *E. Coli*.

25 Figure 5A shows melanin ingestion by keratinocytes treated with SLIGRL, STI or Compound I.

Figure 5B shows the same results as in Figure 5A using isolated melanosomes.

30 Figure 6A shows ICAM-1 immuno-fluorescence staining of treated keratinocytes.

Figure 6B shows a Western blot of immuno-precipitated ICAM-1 protein from treated keratinocytes.

Figure 7A shows human skin, grafted on immuno-suppressed mice, and treated with vehicle or SLIGRL.

Figure 7B shows histological sections of human skin,  
5 grafted on immuno-suppressed mice, and treated with  
vehicle or SLIGRL.

Figure 7C shows histological sections of human skin,  
grafted on immuno-suppressed mice, and treated with  
10 vehicle or STI.

Figure 8 shows scanning electron microscopy images of  
treated keratinocytes.

15 Figure 9 shows F-actin staining of treated  
keratinocytes.

Figure 10 shows the effect of anti-ICAM-1 antibodies on  
keratinocyte phagocytosis.

20

Figure 11 shows the effect of compounds of this  
invention in lightening human age spots.

Detailed Description of the Invention

This invention is based on the discovery that PAR-2-mediated phagocytosis and PAR-2 mediated ICAM-1 expression can be specifically altered. This ability to specifically increase and decrease these cellular functions permits the treatment and prevention of disorders, which would be ameliorated by an increase, or decrease of phagocytosis and/or ICAM-1 expression. Accordingly, this invention provides various compositions and methods for the treatment of disorders ameliorated by the specific alteration of phagocytosis and/or ICAM-1 expression.

More specifically, this invention provides a number of compositions of matter for treating and preventing certain mammalian disorders. These compositions include the following:

(1) a composition of matter for treating a mammal afflicted with a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a therapeutically effective amount of an agent that increases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier;

(2) a composition of matter for treating a mammal afflicted with a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a therapeutically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier;

35



(3) a composition of matter for preventing in a mammal a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a prophylactically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier; and

(4) a composition of matter for preventing in a mammal a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a prophylactically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier.

This invention also provides methods of altering the phagocytosis or ICAM-1 expression level in a cell. The invention first provides a method of increasing phagocytosis or ICAM-1 expression in a mammalian cell, comprising contacting the cell with an effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression. Second, this invention provides a method of decreasing phagocytosis or ICAM-1 expression in a mammalian cell, comprising contacting the cell with an effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.

This invention further provides methods of treatment and prophylaxis regarding disorders affected by the alteration of phagocytosis or ICAM-1 expression. Specifically, this invention provides the following:

(1) a method of treating a mammal afflicted with a disorder ameliorated by an increase in phagocytosis or

ICAM-1 expression in appropriate cells, which comprises administering to the mammal a therapeutically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression;

5

(2) a method of treating a mammal afflicted with a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a therapeutically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression;

(3) a method of preventing in a mammal a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a prophylactically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression; and

(4) a method of preventing in a mammal a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a prophylactically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.

The instant composition of matter can be of any form known in the art. In one embodiment, the composition comprises a pharmaceutically acceptable carrier and one or more discrete pharmaceutical compounds that function as the agent that specifically alters phagocytosis or ICAM-1 expression. In another embodiment, the composition of matter comprises a naturally-occurring composition, or an extract or component thereof, which is deemed pharmaceutically or

cosmetically acceptable. Such naturally occurring compositions contain certain components which function as active agents, and numerous others that serve as pharmaceutical or cosmetically carriers. The instant  
5 compositions can be artificial, naturally occurring, or a combination thereof. In addition, the compositions can be of any physical form known in the art, such as liquids (e.g., solutions, creams, lotions, gels, injectables), solids (e.g., tablets, capsules, powders,  
10 granules), aerosols, and coatings.

Natural compounds that inhibit trypsin, such as serine protease inhibitors, and in particular, soybean trypsin inhibitor ("STI"), can be used for this  
15 invention. Soybean extracts, limabean extracts and similar extracts, and other natural products made from soybean and the like, such as soybean milk, soybean paste, miso, trypsin inhibitor from soybean or limabean and the like, can also reduce phagocytosis by this  
20 mechanism. In the preferred embodiment, the naturally occurring composition is soy milk or STI. Additional sources of serine protease inhibitors include, for example, the following plant families: Solanaceae (e.g., potato, tomato, tomatilla, and the like); Gramineae  
25 (e.g., rice, buckwheat, sorghum, wheat, barley, oats and the like); Cucurbitaceae (e.g., cucumbers, squash, gourd, luffa and the like); and, preferably, Leguminosae (e.g., beans, peas, lentils, peanuts, and the like).

30 As an example, formulations can contain soybean milk or other liquid formulations derived directly from legumes or other suitable plant. In one example, such a formulation contains a large proportion of soybean milk, an emulsifier that maintains th physical stability of  
35 the soybean milk, and optionally, a chelating agent,


preservatives, emollients, humectants and/or thickeners or gelling agents.

The agent in the instant compositions that specifically increases or decreases phagocytosis or ICAM-1 expression can be any type of compound known in the art. Examples include, without limitation, organic molecules, inorganic molecules, peptides, proteins, carbohydrates, nucleic acid molecules, lipids, and any combination thereof. Serine proteases and PAR-2 agonists, for example, can be used to increase phagocytosis. Trypsin, tryptase and thrombin inhibitors and PAR-2 antagonists can be used to decrease phagocytosis.

15

In the preferred embodiment for increasing phagocytosis, the agent is SLIGRL, SAIGRL, or SLIGKVD. In the preferred embodiment for decreasing phagocytosis, the agent is a soybean derivative (such as soybean milk, soybean paste or STI) or Compound I. Compound I has the chemical formula (S)-N-Methyl-D-phenylalanyl-N-[4-[(aminoiminomethyl)amino]-1-(2-benzothiazolylcarbonyl)butyl]-L-prolinamide (as identified in Chemical Abstracts), and has the structure shown below.

25



This compound is described in U.S. Patent No. 5,523,308, as well as in Costanzo, et al., J. Med. Chem., 1996, 39:3039-3043. U.S. Patent No. 5,523,308 describes  
5 related compounds that behave as serine protease inhibitors (such as compounds with a d-phenylalanine-proline-arginine motif), and that can therefore be used to decrease phagocytosis and ICAM-1 expression. Additional compounds related to Compound I are described  
10 in detail in the Examples below.

As used herein, the term "mammal" means any member of the higher vertebrate animals included in the class Mammalia, as defined in Webster's Medical Desk Dictionary  
15 407 (1986), and includes but is not limited to humans, other primates, pigs, dogs, and rodents (such as immunosuppressed mice). In the preferred embodiment of this invention, the mammal is a human.

20 Disorders that can be treated or prevented using the instant invention include any disorder that can be ameliorated (i.e., a positive effect on the disorder per se, and/or its secondary effects) by either an increase or decrease in phagocytosis or ICAM-1  
25 expression in appropriate cells. In the preferred embodiment, the phagocytosis is PAR-2-mediated. These disorders include, without limitation, immune system disorders, diabetes, inflammatory disorders, disorders of the central nervous system, skin disorders, physical  
30 wounds, periodontal disorders and respiratory disorders. These disorders also include, for example, unwanted fertilization, which in one embodiment are prevented by administering inhibitors (i.e. PAR-2 inhibitors) of the sperm protease acrosin which  
35 initiates the PAR-2 pathway (for a discussion of

acrosin, see Fox, et al., FEBS Lett 417:3, 267-9, 1997).

5 A number of disorders have characteristics of more than one category of disorder. Such disorders include, for example, adhesion disorders, which can be categorized as both skin disorders and immune system disorders. Accordingly, a statement herein that a disorder is of a particular category (e.g., skin  
10 disorder) means that, at the very least, the disorder bears traits of that category. Again, however, the disorder may additionally bear traits of another category.

15 Increasing the ability of immune cells to ingest foreign objects like bacteria and viruses would be expected to enhance the immune response. For example, mononuclear phagocytes are inactive in chronic microbial infections (Reiner, Immunol Today 15:8, 374-  
20 81, 1994), and their re-activation would be expected to treat the disease. Alternatively, disorders wherein the immune system is too active would be ameliorated by inhibiting phagocytosis.

25 Immune system and inflammatory disorders treatable in this invention include, by way of example, AIDS, chemotherapy-induced immunodeficiency, asthma, damage due to toxic substance exposure (e.g., asbestos or smoke), host rejection of implants and transplanted  
30 tissue, adhesion disorders, mild infections (such as common colds), severe infections (such as meningitis or "killer bacteria"), wounds (such as infected, diabetic, acute and chronic wounds), restenosis, cystic fibrosis, pulmonary emphysema, periodontal disease, and diaper  
35 rash.

Skin disorders include unwanted pigmentation, unwanted de-pigmentation, psoriasis, rashes, and certain physical skin imperfections (e.g., wrinkles).

5 In one specific example, vitiligo patients are treated with melanin (via liposomes or plain) together with a phagocytosis-increasing agent (e.g., SLIGRL) to darken the light spots. Alternatively, they are treated with Compound I to lighten the darker sites (see U.S. Serial

10 No. 09/110,409, filed July 6, 1998). In an example related to skin disorders, gray hair is treated with melanin (plain or liposome-delivered) and phagocytosis-increasing agent (e.g., SLIGRL), ideally in a shampoo or cream. Central nervous system disorders include,

15 without limitation, Alzheimer's disease and other senile plaque disorders (treated via up-regulating the phagocytosis of amyloid fibrils), depression, phobic disorders, and other disorders resulting from secondary effects of benzodiazepine treatment.

20

The mammalian cells treated in the instant methods are preferably PAR-2-expressing cells, and include, without limitation, keratinocytes, fibroblasts, and "professional phagocytes" (i.e., cells having

25 phagocytosis as a primary function). Professional phagocytes include, for example, neutrophils, macrophages and macrophage-like cells (e.g., Langerhans cells and Kupfer cells). In the preferred embodiment, the mammalian cells are human cells.

30

In this invention, the "appropriate cells" in which phagocytosis or ICAM-1 expression must be altered in response to the instant compositions of matter are readily determined based on the nature of the disorder

35 being treated or prevented. For example, if the

disorder being treated is a pigmentation disorder, the appropriate cells in which phagocytosis or ICAM-1 expression needs to be altered are keratinocytes.

5           The instant methods are directed at preventing as well as treating disorders. As used herein, "treating" a disorder means reducing the disorder's progression, ceasing the disorder's progression, ceasing or otherwise ameliorating secondary effects of the  
10 disorder, reversing the disorder's progression, or preferably, curing the disorder. As used herein, "preventing" a disorder means reducing, and preferably eliminating, the likelihood of the disorder's  
15 occurrence.

15

          In this invention, administering the instant compositions can be effected or performed using any of the various methods and delivery systems known to those skilled in the art. The administering can be performed,  
20 for example, intravenously, orally, via implant, transmucosally, topically, transdermally, intramuscularly, subcutaneously, and via aerosol. In addition, the instant compositions ideally contain one or more routinely used pharmaceutically or cosmetically  
25 acceptable carriers. Such carriers are well known to those skilled in the art. The following delivery systems, which employ a number of routinely used carriers, are only representative of the many embodiments envisioned for administering the instant  
30 composition.

          Transdermal delivery systems include patches, gels, tapes, lotions, soaps, shampoos and creams, and can contain excipients such as solubilizers, permeation  
35 enhancers (e.g., fatty acids, fatty acid esters, fatty



alcohols and amino acids), hydrophilic polymers ( .g., polycarbophil and polyvinylpyrrolidone), and adhesives and tackifiers (e.g., polyisobutylenes, silicone-based adhesives, acrylates and polybutene).

5

Topical delivery of some of the compositions of this invention, particularly those comprising proteins such as trypsin, tryptase and STI, can be achieved using liposomes. The liposomes are preferably non-  
10 ionic. In one example, they contain (a) glycerol dilaurate; (b) compounds having the steroid backbone found in cholesterol; and (c) fatty acid ethers having from about 12 to about 18 carbon atoms, wherein the constituent compounds of the liposomes are in a ratio  
15 of about 37.5:12.5:33.3:16.7. Liposomes comprising glycerol dilaurate/cholesterol/ polyoxyethylene-10-stearyl ether/polyoxyethylene-9-lauryl ether ("GDL" liposomes) are preferred. In one embodiment, the liposomes are present in an amount, based upon the  
20 total volume of the composition, of from about 10 mg/ml to about 100 mg/ml, and preferably from about 15 mg/ml to about 50 mg/ml. A ratio of about 37.5:12.5:33.3:16.7 is preferred. Methods of preparing liposomes are well known in the art, such as those disclosed in  
25 Niemiec, et al., 12 Pharm. Res. 1184-88 (1995).

Also, for topical or transdermal administration, the instant compositions can be combined with other components such as moisturizers, cosmetic adjuvants,  
30 anti-oxidants, bleaching agents, tyrosinase inhibitors and other known depigmentation agents, alpha-hydroxy acids, surfactants, foaming agents, conditioners, humectants, fragrances, viscosifiers, buffering agents, preservatives, sunscreens and the like. The  
35 compositions of this invention can also contain active

amounts of retinoids including, for example, tretinoin, retinol, esters of tretinoin and/or retinol and the like.

5 Transmucosal delivery systems include patches, tablets, suppositories, pessaries, gels and creams, and can contain excipients such as solubilizers and enhancers (e.g., propylene glycol, bile salts and amino acids), and other vehicles (e.g., polyethylene glycol, 10 fatty acid esters and derivatives, and hydrophilic polymers such as hydroxypropylmethylcellulose and hyaluronic acid).

Injectable drug delivery systems include 15 solutions, suspensions, gels, microspheres and polymeric injectables, and can comprise excipients such as solubility-altering agents (e.g., ethanol, propylene glycol and sucrose) and polymers (e.g., polycaprylactones and PLGA's). Systems for central 20 nervous system delivery include, for example, a lipid-coupled derivative to cross the blood brain barrier (e.g. DHA). Implantable systems include rods and discs, and can contain excipients such as PLGA and polycaprylactone.

25

Oral delivery systems include tablets and capsules. These can contain excipients such as binders (e.g., hydroxypropylmethylcellulose, polyvinyl pyrrolidone, other cellulosic materials and starch), 30 diluents (e.g., lactose and other sugars, starch, dicalcium phosphate and cellulosic materials), disintegrating agents (e.g., starch polymers and cellulosic materials) and lubricating agents (e.g., stearates and talc). Such delivery systems also

include, for example, toothpaste, mouthwash, lozenges and lollipops.

Solutions, suspensions and powders for  
5 reconstitutable delivery systems include vehicles such as suspending agents (e.g., gums, zanthans, cellulosics and sugars), humectants (e.g., sorbitol), solubilizers (e.g., ethanol, water, PEG and propylene glycol),  
10 surfactants (e.g., sodium lauryl sulfate, Spans, Tweens, and cetyl pyridine), preservatives and antioxidants (e.g., parabens, vitamins E and C, ascorbic acid, and natural extracts), anti-caking agents, coating agents, and chelating agents (e.g., EDTA). Oil-in-water emulsions, water-in-oil emulsions,  
15 solvent-based formulations and aqueous gels known to those of skill in the art can also be utilized as vehicles for the delivery of the compositions of this invention.

20 Methods of determining therapeutically and prophylactically effective doses for administering the instant compositions in humans are known in the art. For example, these effective doses can readily be determined from the results of animal studies.

25

In one example, the instant composition is applied to the skin surface such that, based upon a square cm of skin surface, from about 2  $\mu\text{l}/\text{cm}^2$  to about 200  $\mu\text{l}/\text{cm}^2$  of phagocytosis-altering agent is present when a change  
30 in phagocytosis is desired. When using a thrombin and trypsin inhibitor such as Compound I or its analogs, whether synthetically- or naturally-derived in a formulation, such an active compound is present in an amount of from about 0.0001% to about 15% by  
35 weight/volum of the composition. In an ther

embodiment, it is present in an amount of from about 0.0005% to about 5% of the composition. Preferably, it is present in an amount of from about 0.001 to about 1% of the composition.

5

In another example, liquid derivatives and natural extracts made directly from plants or botanical sources are employed in the instant compositions in a concentration (w/v) of from about 1 to about 99%, and preferably from about 75 to about 95%. In still another example, fractions of natural extracts and naturally derived protease inhibitors such as STI have a concentration range of from about 0.01% to about 20% and, preferably, from about 1% to about 10% of the composition.

10  
15

This invention still further provides an article of manufacture for administering to a mammal the instant composition of matter, comprising a solid delivery vehicle having the composition operably (i.e., deliverably) affixed thereto. The solid delivery vehicle can be any device designed to come into temporary or permanent contact with the body, whether or not it was originally intended for use as a delivery vehicle. Examples of the instant article of manufacture include, without limitation, coated bandages or other wound dressing for treating wounds, coated bodily implants (including implants with coated internal scaffolding) for either preventing or promoting tissue growth, and coated balloon catheters and stents for preventing restenosis.

Finally, this invention provides a method of administering a therapeutic, prophylactic or cosmetic compound to a mammal, comprising administering to the

20  
25  
30  
35

mammal (a) the compound and (b) a composition of matter comprising a pharmaceutical or cosmetic carrier and an agent that specifically increases phagocytosis in an amount sufficient to increase phagocytosis in cells  
5 where uptake of the compound is desired, wherein the composition is administered prior to and/or concurrently with the administration of the compound.

The pharmaceutical compound can be, for example, a  
10 polypeptide, protein, or nucleic acid molecule. In one embodiment, the pharmaceutical compound and composition are administered together via microscopic porous biodegradable beads, which then release the pharmaceutical compound after being ingested through  
15 phagocytosis by the appropriate cells.

This invention will be better understood by reference to the Examples which follow, but those skilled in the art will readily appreciate that they are only  
20 illustrative of the invention as described more fully in the claims which follow thereafter. In addition, various documents are cited throughout this application. The disclosures of these documents are hereby incorporated by reference into this application to describe more fully  
25 the state of the art to which this invention pertains.

Examples5            Example 1  
             SLIGRL, STI and Compound I Affect Keratinocyte  
             Phagocytosis

             In order to study the role of the PAR-2 pathway in  
phagocytosis, several *in vitro* model systems were used.  
10 One system used contained primary human keratinocytes or  
a human keratinocyte cell line. In this and a number of  
following examples, cells were treated with test  
compounds for different amounts of time (from one hour to  
three days), and samples were then incubated with  
15 fluorescent microspheres for two hours. The beads  
ingested by the cells were photographed using  
fluorescence microscopy.

             In this example, human primary keratinocytes or the  
20 human keratinocyte cell line HaCaT were used as *in vitro*  
model systems to study the effect of PAR-2 regulators on  
keratinocyte phagocytosis. The human primary  
keratinocytes used are commercially available from  
Clonetics (San Diego, CA). Cells were plated on chamber  
25 slides, at 2 chambers/slide and 60,000 cells/chamber.  
Cells were treated once daily, for two or three days,  
with Compound I (1 $\mu$ M), SLIGRL (10 $\mu$ M), or vehicle  
(Phosphate-buffered saline, ("PBS") from Gibco-BRL  
(Gaithersburg, MD). After two or three days of exposure  
30 to the test compounds, cells were exposed to Nile-red or  
FITC fluorescent microspheres, 1  $\mu$ m in diameter, 50  
microspheres/cell, for two hours at 37°C. Microspheres  
were from Molecular Probes (Eugene, OR), and were  
processed according to manufacturer's instructions.  
35 Following that treatment, cells were incubated with 15%  
Fetal Bovine Serum ("FBS", from Gibco-BRL), for 15

minutes at 37°C and rinsed with PBS. At that time, chambers were separated from the slides, and the slides were covered with glycerol and coverslips. Fluorescent microscopy was performed using a Zeiss Axiovert 35 or a  
5 Nikon Optiphot-2 microscope.

Figure 1 shows three images of human primary keratinocytes, treated for two days with vehicle (control), Compound I or SLIGRL. As seen in this Figure,  
10 the microspheres were ingested by the control keratinocytes, and were distributed around the cell's nucleus. Microspheres were also found around the cell, probably because they are non-specifically attached to extracellular-matrix components secreted by the cell.  
15 The quantity of the microspheres ingested was changed with the treatments. Treatment with Compound I, an inhibitor of PAR-2 activation results in a dramatic reduction in the quantity of ingested microspheres. Treatment with SLIGRL, a PAR-2-activating peptide,  
20 results in a dramatic increase in the number of ingested microspheres.

The same results were also obtained when the human keratinocyte cell line HaCaT was used instead of the  
25 primary keratinocytes (see Figure 2). SLIGRL, STI and Compound I were tested for their effect on keratinocyte phagocytosis. In these experiments, the extracellular accumulation of microspheres could be washed off. The only particles visible were microspheres internalized by  
30 the keratinocytes, which were accumulated around their nuclei. Soybean trypsin inhibitor ("STI"), which is a serine protease inhibitor capable of affecting the PAR-2 pathway, was shown to reduce microsphere ingestion in this experiment, as was Compound I. SLIGRL treatment,  
35 on the other hand, resulted in increased microsphere

ingestion. Each of these experiments was repeated at least three times. These experiments show, for the first time, that keratinocytes have PAR-2-mediated phagocytic ability. These experiments also demonstrate that  
5 compounds that regulate the PAR-2 pathway can regulate the level of keratinocyte phagocytosis.

When these experiments were repeated using melanocytes, which do not express PAR-2, SLIGRL had no  
10 inducible effect on microsphere ingestion. Melanocytes did not ingest beads under any of the above conditions. Since SLIGRL activates PAR-2 only, and melanocytes do not express PAR-2, these cells do not respond to the SLIGRL signal and phagocytosis cannot be affected.

15

Example 2  
SLIGRL and Compound I Affect Fibroblast Phagocytosis

The experiment described in Example 1 was repeated  
20 using the fibroblast cell line 92-3T3 (obtained from the ATCC in Rockville, MD). Fibroblasts are not known to possess phagocytic ability. Indeed, only minimal bead ingestion was observed with untreated fibroblasts. However, SLIGRL-treated fibroblasts increased the  
25 number of ingested beads (Figure 3). SLIGRL-induced fibroblast phagocytosis was quantitatively different from that of keratinocytes, since fibroblasts do not perform phagocytic tasks in vivo. This experiment shows, for the first time, that fibroblasts have  
30 inducible PAR-2 phagocytic ability. In other words, this experiment demonstrates that compounds that regulate the PAR-2 pathway can regulate the level of fibroblast phagocytosis.



Example 3SLIGRL, STI and Compound I Aff ct Macrophage Phagocytosis

5           The experiment described in Example 1 was repeated using the macrophage cell line IC-21 (obtained from the ATCC), which shares phagocytic characteristics with peritoneal macrophages. As shown for keratinocytes and fibroblasts, Compound I and STI reduced, and SLIGRL  
10 increased, the number of microspheres ingested by these macrophages which are "professional phagocytic" cells.

To better quantify the level of phagocytosis, the "Vybrant™ Phagocytosis Assay Kit" of Molecular Probes  
15 (Eugene, Oregon) was used, following manufacturer's instructions, with modification of cell culture conditions for the IC-21 cell line. This kit uses Fluorescein-labeled *E. Coli* K-12 particles, and is designed for quantifying the effects of drugs or other  
20 environmental factors on phagocytic functions. Macrophages were treated overnight with 100nM of Compound I, 5μM of SLIGRL, or 0.1 mg/ml of STI, all dissolved in PBS. The ability of the treated macrophages to ingest the fluorescent *E. Coli*, as  
25 measured by this kit, is documented in Table 1. This experiment was repeated three times. Table 1 represents data from one experiment.

Table 1

<u>Treatment</u>	<u>% effect (ingestion)</u>
Untreated control	100
SLIGRL	331.6 +/- 5.9
Compound I	89.9 +/- 13.6
STI	56.06 +/- 12.4

35

This experiment demonstrates that macrophage phagocytosis can be regulated by PAR-2 pathway modulators. It also shows that both synthetic compounds and naturally derived compounds can modulate phagocytosis via the PAR-2 pathway.

Example 4

Dose-response Relationship Between PAR-2 Signaling and Macrophage Phagocytosis

10

In order to verify the quantitative nature of the macrophage-phagocytosis assay, a dose-response experiment was performed. Macrophages were treated with 0, 0.01, 0.1 and 1 mg/ml of STI, and the experiment was performed as described in Example 3. A dose-response of decreased phagocytosis with increasing STI concentrations was observed, as indicated in Figure 4A. Similar results were obtained for Compound I at 0.01, 0.1 and 1 nM, while SLIGRL treatment resulted in an increase in phagocytosis (Figure 4B). Each experiment was repeated three times. This experiment demonstrates that the phagocytic effect of PAR-2-modulating compounds is dose-responsive and can be quantified.

25

Example 5

Dose-response Relationship Between PAR-2 Signaling and Keratinocyte Phagocytosis

Human keratinocytes were treated with increasing concentrations of SLIGRL, the PAR-2 peptide activator and agonist, at 0, 5 and 10  $\mu$ M for two days in the same manner as set forth in Example 1. Increasing concentrations of SLIGRL result in increased phagocytosis. Human keratinocytes were also treated with increasing concentrations of Compound I and STI for two days. Treatment with increasing concentrations of

Compound I (from 1pM to 1 $\mu$ M) or with STI (from 0.01 to 1 mg/ml), results in a dose-dependent decrease in phagocytosis (see Table 2).

5 Image analysis of the fluorescent beads inside the keratinocytes was used as an alternative way to quantify the phagocytic effect in this system. Empire Imagins Database Version 1.1 was used on a Gateway 2000 P5-100 computer (Media Cybernetics, Silver Springs, MD) for  
 10 capturing images. Image Pro Plus version 1.3 was used for measurements, and Microsoft Excel version 5.0 was used for data processing. Data obtained from this keratinocyte-microsphere system were in full agreement with data from the macrophage/*E. Coli* system.

15

Table 2

	<u>Treatment</u>	<u>% Ingestion</u>
	Untreated	100 +/- 12
	STI, 0.01%	76 +/- 15
20	STI, 0.1%	55 +/- 14
	STI, 1%	41.6 +/- 11

Example 6

25 Compound I, SLIGRL and STI Affect the Acquisition of Pigment by Keratinocytes

This example tests the ability of keratinocytes to acquire melanin or melanosomes *in vitro*, and thus function as a simplified system for melanosome uptake in  
 30 the skin. Keratinocytes were plated in glass chamber-slides as described earlier, and were treated for two days with SLIGRL, Compound I or STI. At that time, melanin powder (from Sigma, St. Louis, MO) was mixed in sterile PBS at 10  $\mu$ g/ml, and was added to the culture  
 35 media (1:10 dilution) for two hours. Cells were then washed with PBS and stained with Fontana-Mason ("F&M")

staining. F&M stains silver nitrate-reducing molecules, thereby permitting the identification of melanins inside the keratinocytes. As shown in Figure 5A, untreated keratinocytes were able to ingest melanin from the culture media, and localize the internalized melanin around their nuclei. This system, therefore, can mimic melanosome transfer and melanin distribution *in vivo*, as skin keratinocytes use melanin as an UV-protective cap over their nuclei. This capping pattern is also observed with the ingested microspheres as demonstrated in Example 1, and Figures 1 and 2. Figure 5A also shows that the SLIGRL treatment, which turns on the PAR-2 pathway, dramatically increases the internalization of melanin and its deposition around the nuclei. Compound I and STI, on the other hand, dramatically reduce the uptake of melanin by the keratinocytes. This example demonstrates that PAR-2-modulating agents, of both synthetic and natural origin, can affect pigment distribution in epidermal cells. The same results were also observed using melanosomes isolated as described in S. Orlow, et al., J.I.D. 100:55-64 (1993) (Figure 5B).

Example 7  
Cell-Cell Contact is Required for Compound I Effect on Pigment Transfer from Melanocytes to Keratinocytes.

Since PAR-2 is expressed in keratinocytes, but not in melanocytes, the possible requirement of keratinocyte-melanocyte contact was tested for the effect of Compound I and SLIGRL on melanosome phagocytosis by the keratinocytes. Primary melanocyte cultures (commercially available from Clonetics, San Diego) were plated under epidermal equivalents (EpiDerm, of MatTek, Ashland, MA) to create an equivalent-monolayer co-culture with no contact between keratinocytes and melanocytes. These co-cultures were compared to MelanoDerm equivalents (of

MatTek), where melanocytes are present in the basal layer of the equivalent. Cultures were treated with Compound I, with the PAR-2 agonist TFLLRNPNDK, and with the PAR-2 agonist SLIGRL. As set forth in Table 3, keratinocytes are indicated by "K", melanocytes are indicated by "M", and lack of keratinocyte-melanocyte contact is indicated as "no K-M cont". As shown in Table 3, no effect on melanosome transfer was observed in equivalent-monolayer co-cultures (having no keratinocyte/ melanocyte contact) treated with these agents, when measured as the level of pigmentation. In melanocyte-containing equivalents, Compound I reduced and SLIGRL induced pigmentation by affecting melanosome transfer. The same result was also observed with monolayer keratinocyte/melanocyte co-cultures having keratinocyte/melanocyte contact. These results demonstrate that keratinocyte-melanocyte contact is required for the PAR-2 effect on melanosome phagocytosis.

20

Table 3

25

<u>Treatment</u>	<u>Mono-Layer Co-Cultures</u>	<u>Equivalent Monolayer Co-culture</u>	<u>Epidermal Equivalents</u>
	(K-M cont.)	(no K-M cont.)	(K-M cont.)

Compound I	lightening	no effect	lightening
SLIGRL	darkening	no effect	darkening
TFLLRNPNDK	no effect	no effect	no effect

30

Example 8  
Timing of the Effect of Compound I and SLIGRL on Phagocytosis

Human keratinocytes were treated with Compound I or SLIGRL for periods of time ranging from one hour to two days. At the end of the treatment period, the cells were

treated with microspheres as described in Example 1. Table 4 shows the time required for these compounds to affect phagocytosis. Plus signs indicate an effect on phagocytosis, minus signs indicate no effect on phagocytosis, and plus/minus signs indicate a marginal effect on phagocytosis. The effects measured were a decrease for Compound I and an increase for SLIGRL. This experiment demonstrates that following the activation or the inhibition of the PAR-2-signaling pathway, at least eight hours are required to alter the phagocytic ability of the keratinocytes. This implies that the PAR-2 signaling results in new protein synthesis, a reduction in protein synthesis when a turnover time is required to eliminate the existing relevant protein(s), or a rearrangement of proteins (as occurs in the reorganization of cytoskeletal components).

Table 4

	<u>Time of Treatment</u>	<u>Compound I</u>	<u>SLIGRL</u>
20	1 hour	-	-
	2 hours	-	-
	4 hours	-	-
	6 hours	-	-
25	8 hours	-/+	-/+
	16 hours	+	+
	24 hours	+	+
	48 hours	+	+

30 Example 9  
Compound I, STI and SLIGRL Affect ICAM-1  
Intracellular Expression and Localization

Intercellular adhesion molecule-1 (ICAM-1) is an inducible cell-surface glycoprotein that is implicated in cell-cell adhesion, cell membrane ruffling and

phagocytosis. Therefore, the effect of PAR-2 modulation on ICAM-1 was tested. Keratinocytes were grown in chamber slides and treated with SLIGRL, Compound I and STI as described, followed by immunofluorescent staining for ICAM-1 using standard procedures. Normal donkey serum, (used at 1:5 dilution), was obtained from Jackson ImmunoResearch Laboratory (Westgrove, PA). A polyclonal goat anti-human ICAM-1 antibody, (used at 1:200 dilution), was obtained from R&D Systems (Minneapolis, MN), and FITC-conjugated donkey anti-goat antibody was obtained from Jackson ImmunoResearch Laboratory. As shown in Figure 6A, the intracellular localization of ICAM-1 can be modulated via the PAR-2 pathway. In untreated keratinocytes, ICAM-1 was localized mainly to the plasma membrane, with some staining at the nuclear membrane. Following Compound I or STI treatment, less staining was observed at the cytoplasmic membrane, and more at the nuclear membrane. SLIGRL treatment had the opposite effect, resulting in increased and more diffused cytoplasmic membrane staining, and reduced nuclear membrane staining.

Immuno-precipitation and Western blotting experiments were performed, according to known methods, to assess the level of ICAM-1 protein in treated keratinocytes. As shown in Figure 6B, SLIGRL increased the level of ICAM-1 expression in these cells, whereas Compound I decreased the level of ICAM-1 expression.

30

Example 10  
STI and Compound I Affect Chemotactic Cell Migration

35 Since ICAM-1 is involved in cell migration, cell migration towards a chemotactic peptide was studied in

cells treated with Compound I and STI. Human PMN cells were placed in on side of a Boyden chamber, using standard techniques, and a chemotactic peptide (FMLP) was placed in the other chamber. Cells were allowed to

5 migrate into the second chamber, and the number of migrating cells per field and distance of migration were calculated. The experiment was repeated with PMN cells pretreated with one of the following: 5 or 0.5 mg/ml STI, 1 or 0.1  $\mu$ M Compound I, a buffer vehicle,

10 and with and without chemotactic peptide. The number of cells per field that migrated the same distance as the untreated control was measured. These data are summarized in Table 5. Without the peptide and with no treatment, an average of 19 cells/field were migrating

15 a distance of 80 microns. The addition of the chemotactic peptide resulted in 41 cells/field migrating 115 microns. Both compounds completely inhibited the migration of cells towards this peptide. In other words, no cells (or less than 5/field) were

20 identified at 115 microns. This indicates that these inhibitors affect not only phagocytosis, but also cell migration. The ability to inhibit PMN migration is an important constituent of anti-inflammatory compounds.

25

Table 5

<u>Treatment</u>	<u>Cells/Field</u>	<u>Migration</u>
Buffer	19	80 $\mu$
FMLP	41	115
FMLP + Cpd I	<5	115
30 FMLP+ STI	<5	115



Example 11  
SLIGRL and STI Affect Human Skin Pigmentation

5 Human white facial skin samples were grafted on immuno-suppressed mice using standard techniques. About six weeks later grafts of the same individual, grafted on different mice, were treated with vehicle (ethanol: propylene glycol 70:30), or with 50  $\mu$ M  
10 SLIGRL. Treatment was performed daily, 5 days/week. Darkening of the SLIGRL-treated grafts was visually observed during the last weeks of treatment (see Figure 7A). On day 66, the animals were sacrificed and their skins were analyzed histologically. F&M-stained  
15 sections revealed an increase in melanin in the SLIGRL-treated grafts, as shown in Figure 7B. This experiment demonstrates the ability to use SLIGRL on human skin to induce sunless tanning.

20 The same experiment was repeated with black human breast skin samples grafted on immuno-suppressed mice. Grafts of the same individual were treated with vehicle (GDL liposomes) or with 1% STI. GDL liposomes were prepared as set forth in Niemiec, et al., with the  
25 exception of the following changes: the non-ionic liposomal formulation contained glycerol dilaurate (Emulsynt GDL, ISP Van Dyk)/cholesterol (Croda)/polyoxyethylene-10-stearyl ether (Brij76, ICI)/polyoxyethylene-9-lauryl ether, at ratio of  
30 37.5:12.5:33.3:16.7. Hepes buffer, 0.05M, pH 7.4 (Gibco-BRL of Gaithersburg, MD) was used as the aqueous phase in the preparation of the liposomes. Figure 7C shows F&M-stained skin sections from these grafts. This figure clearly demonstrates the ability of STI to reduce  
35 pigmentation in human skin.

Example 12PAR-2 Effect on Phagocytosis and Migration Relates to Changes in Cell Shape

5

Both phagocytosis and cell migration involve changes in cell shape. Therefore, the effect of the PAR-2 pathway reagents on cell podia were studied using scanning electron microscopy. Keratinocytes were treated with Compound I (identified as "SH00230"), SLIGRL and a buffer vehicle for two days, and then processed for SEM using standard techniques. As shown in Figure 8, a dramatic change in the shape of cell podia is observed. Relative to the vehicle-treated controls, the Compound I-treated cells have dramatically shorter podia. In other words, the cells' "fingers" are shorter and malformed, so they cannot grasp objects as well as can the control cells. The SLIGRL-treated samples demonstrated the opposite effect. Their podia were increased in number, were somewhat longer, and were much thinner. In other words, these cells have a greater probability for productive interaction with particles. Comparing these SEM pictures in Figure 8, it is clear that SLIGRL-treated cells have a greater ability to interact with particles, while Compound I-treated cells are less suited for such a task.

30 Example 13PAR-2 Affects Cytoskeletal Organization

Changes in cell shape as demonstrated in Figure 8 require reorganization of cytoskeletal components. Therefore, the organization of F-actin filaments following PAR-2 modulation was tested. Keratinocytes were treated with Compound I (10 nM), STI (0.1 mg/ml), or with SLIGRL (5  $\mu$ M) as described in Example 1, and

stained for F-actin using standard techniques. Figure 9 shows dramatic changes in actin filaments organization following these treatments. SLIGRL treatment induced actin polymerization around the cell cortex, an area important in the control of cell movement and phagocytosis. STI and Compound I, on the contrary, reduced the ordered organization of the cell cortex, thereby reducing the cells' ability to regulate their movement and podia.

10

Example 14  
ICAM-1 Modulation Affects Keratinocyte  
Phagocytosis

15 Keratinocytes were exposed to fluorescent microspheres (i.e. beads) as described in Example 1. These cells were pretreated with 50 µg/ml mouse anti-human ICAM-1 antibodies (from R&D Systems) for 16 hours, and then boosted again for four hours before  
20 incubating with the beads. As shown in Figure 10, blocking the surface ICAM-1 molecules on the keratinocytes results in reduced bead ingestion. This experiment therefore establishes a link between ICAM-1 and keratinocyte phagocytosis.

25

Example 15  
STI/Liposome Formulation Can Lighten Human Age  
Spots

30 An individual with three age spots on the dorsum of her hand was treated for eight weeks, twice a day, as follows. The proximal age spot was treated with placebo, containing 20 mg/ml of liposomes. The median age spot was not treated. The distal age spot was  
35 treated with STI, 1%, in liposomes (20 mg/ml). GDL liposomes were prepared as set forth in Niemiec, et al., with the exception of the following changes. The non-ionic liposomal formulation contained glycerol

dilaurate (Emulsynt GDL, ISP Van Dyk)/cholesterol (Croda)/polyoxyethylene-10-stearyl ether (Brij76, ICI)/polyoxyethylene-9-lauryl ether, as at ratio of 37.5:12.5:33.3:16.7. Hepes buffer, 0.05M, pH 7.4

5 (Gibco-BRL of Gaithersburg, MD) was used as the aqueous phase in the preparation of the liposomes. UV and visible light digital pictures were taken at times of 0, 4 and 8 weeks of treatment. L\* (brightness) values were calculated from the images using Adobe Photoshop.

10

As shown in Figure 11, the age spot treated with STI became lighter following 8 weeks of treatment. Figure 11 is a composite of four pictures. The left panel is a visible light picture of the hand, before

15 (upper) and after (lower) 8 weeks of treatment. At this orientation, the proximal age spot is placebo-treated, the median age spot is untreated, and the distal age spot is STI-treated. The right panel shows the same hand at the same time points, using UV-

20 photography. UV light enables the visualization of pigment deeper in the skin, demonstrating that the STI whitening effect is not superficial. Figure 11 clearly demonstrates that the STI formulation lightened the distal age-spot. An increase of 15 L\* units was


25 calculated for this STI-treated spot, further demonstrating the ability of this treatment to lighten age spots.

30 Example 16  
Phagocytosis-Reducing Compounds Analogous to Compound I

Certain compounds, and their pharmaceutically acceptable salts, such as those described in Costanzo, et al., "Potent Thrombin Inhibitors That Probe the S<sub>1</sub>' Subsite: Triptide Transition State Analogues Based on a Het rocycl -Activated Carbonyl Group", J. Med. Chem.,

1996, Vol. 39, pp. 3039-3043, behave as serine protease inhibitors (i.e. phagocytosis inhibitors), and have the following structural formula:

5



10

wherein:

15

A is selected from the group consisting of C<sub>1</sub>-  
 8alkyl, carboxyC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxycarbonylC<sub>1</sub>-4alkyl,  
 phenylC<sub>1</sub>-4alkyl, substituted phenylC<sub>1</sub>-4alkyl (where the  
 phenyl substituents are independently selected from one  
 20 or more of, C<sub>1</sub>-4 alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy,  
 hydroxy, halo, amido, nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-  
 4dialkylamino, carboxy or C<sub>1</sub>-4 alkoxycarbonyl), formyl,  
 C<sub>1</sub>-4alkoxycarbonyl, C<sub>1</sub>-2alkylcarbonyl, phenylC<sub>1</sub>-  
 4alkoxycarbonyl, C<sub>3</sub>-7cycloalkylcarbonyl,  
 25 phenylcarbonyl, substituted phenylcarbonyl (where the  
 phenyl substituents are independently selected from  
 one or more of C<sub>1</sub>-4alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-  
 4alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1</sub>-  
 4alkylamino, C<sub>1</sub>-4dialkylamino, carboxy or C<sub>1</sub>-4  
 30 alkoxycarbonyl), C<sub>1</sub>-4alkylsulfonyl, C<sub>1</sub>-4alkoxysulfonyl,  
 perfluoroC<sub>1</sub>-4alkyl-sulfonyl, phenylsulfonyl,  
 substituted phenylsulfonyl (where the phenyl  
 substituents are independently selected from one or  
 more of C<sub>1</sub>-4alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy,  
 35 hydroxy, halo, amido, nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-  
 4dialkylamino, carboxy or C<sub>1</sub>-4 alkoxycarbonyl), 10-  
 camphorsulfonyl, phenylC<sub>1</sub>-4alkylsulfonyl, substituted  
 phenylC<sub>1</sub>-4alkylsulfonyl, C<sub>1</sub>-4alkylsulfinyl, perfluoroC<sub>1</sub>-

4alkylsulfinyl, phenylsulfinyl, substituted phenylsulfinyl (where the phenyl substituents are independently selected from one or more of, C<sub>1</sub>-4alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy, hydroxy, halo, amido, 5 nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-4dialkylamino, carboxy or C<sub>1</sub>-4alkoxycarbonyl), phenylC<sub>1</sub>-4alkylsulfinyl, substituted phenylC<sub>1</sub>-4alkylsulfinyl, 1-naphthylsulfonyl, 2-naphthylsulfonyl or substituted naphthylsulfonyl (where the naphthyl substituents are 10 independently selected from one or more of, C<sub>1</sub>-4alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy, hydroxy, halo, amido, nitro, amino, carboxy or C<sub>1</sub>-4alkoxycarbonyl), 1-naphthylsulfinyl, 2-naphthylsulfinyl or substituted naphthylsulfinyl (where the naphthyl substituents are 15 independently selected from one or more of, C<sub>1</sub>-4alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-4dialkylamino, carboxy or C<sub>1</sub>-4alkoxycarbonyl);

20 a D or L amino acid which is coupled as its carboxy terminus to the nitrogen depicted in the structure above and is selected from the group consisting of alanine, asparagine, 2-azetidincarboxylic acid, glycine, N-C<sub>1</sub>-8alkylglycine, proline, 1- 25 amino-1-cycloC<sub>3</sub>-8alkylcarboxylic acid, thiazolidine-4-carboxylic acid, 5,5-dimethylthiazolidine-4-carboxylic acid, oxazolidine-4-carboxylic acid, pipercolinic acid, valine, methionine, cysteine, serine, threonine, norleucine, leucine, tert-leucine, isoleucine, 30 phenylalanine, 1-naphthalanine, 2-naphthalamine, 2-thienylalanine, 3-thienylalanine, [1,2,3,4]-tetrahydroisoquinoline-1-carboxylic acid and [1,2,3,4,]-tetrahydroisoquinoline-2-carboxylic acid

where the amino terminus of said amino acid is connected to a member selected from the group consisting of C<sub>1-4</sub>alkyl, tetrazol-5yl-C<sub>1-2</sub>alkyl, carboxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxycarbonylC<sub>1-4</sub>alkyl, 5 phenylC<sub>1-4</sub>alkyl, substituted phenyl C<sub>1-4</sub> alkyl (where the phenyl substituents are independently selected from one or more of, C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxyl or C<sub>1-4</sub>alkoxycarbonyl), 1,1-diphenylC<sub>1-4</sub>alkyl, 3-phenyl-2-10 hydroxypropionyl, 2,2-diphenyl-1-hydroxyethylcarbonyl, [1,2,3,4]-tetrahydroisoquinoline-1-carbonyl, [1,2,3,4]-tetrahydroisoquinoline-3-carbonyl, 1-methylamino-1-cyclohexanecarbonyl, 1-hydroxy-1-cyclohexanecarbonyl, 15 1-hydroxy-1-phenylacetyl, 1-cyclohexyl-1-hydroxyacetyl, 3-phenyl-2-hydroxypropionyl, 3,3-diphenyl-2-hydroxypropionyl, 3-cyclohexyl-2-hydroxypropionyl, formyl, C<sub>1-4</sub>alkoxycarbonyl, C<sub>1-12</sub>alkylcarbonyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkylcarbonyl, phenylC<sub>1-20</sub> alkylcarbonyl, substituted phenylC<sub>1-4</sub>alkylcarbonyl (where the phenyl substituents are independently selected from one or more of, C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxy or C<sub>1-25</sub> alkoxycarbonyl) 1,1-diphenylC<sub>1-4</sub>alkylcarbonyl, substituted 1,1-diphenylC<sub>1-4</sub>alkylcarbonyl (where the phenyl substituents are independently selected from one or more of, C<sub>1-4</sub>alkyl, perfluoro C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-30</sub> dialkylamino, carboxy or C<sub>1-4</sub> alkoxy-carbonyl), perfluoroC<sub>1-4</sub>alkylsulfonyl, C<sub>1-4</sub>alkylsulfonyl, C<sub>1-4</sub>alkoxysulfonyl, phenylsulfonyl, substituted phenylsulfonyl (where the phenyl substituents are



independently selected from one or more of, C-1alkyl, perfluoroC<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxyl or C<sub>1-4</sub>alkoxycarbonyl), 10-camphorsulfonyl, phenylC<sub>1-4</sub>alkylsulfonyl, substituted phenylC<sub>1-4</sub>alkylsulfonyl, 5 perfluoroC<sub>1-4</sub>alkylsulfinyl, C-1-4alkylsulfinyl, phenylsulfinyl, substituted phenylsulfinyl (where the phenyl substituents are independently selected from one or more of, C<sub>1-4</sub>alkyl, perfluoro C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub> dialkylamino, carboxy or C<sub>1-4</sub>alkoxycarbonyl), 1-naphthylsulfonyl, 1,2-naphthylsulfonyl, substituted naphthylsulfonyl (where the naphthyl substituents are independently selected from one or more of, C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxyl or C<sub>1-4</sub>alkoxycarbonyl), 1-naphthylsulfinyl, 2-naphthylsulfinyl, and substituted naphthylsulfinyl (where the naphthyl substituents are independently selected from one or more of, C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxy or C<sub>1-4</sub>alkoxycarbonyl);

or a polypeptide comprising two amino acids,

25

wherein the first amino acid is a D or L amino acid, bound via its carboxy terminus to the nitrogen depicted in Formula I and is selected from the group consisting of glycine, N-C<sub>1-8</sub>alkylglycine, alanine, 2-azetidinedicarboxylic acid, proline, thiazolidine-4-carboxylic acid, 5,5-dimethylthiazolidine-4-carboxylic acid, oxazolidine-4-carboxylic acid, 1-amino-1-cycloC<sub>3-8</sub>alkylcarboxylic acid, 3-hydroxyproline, 4-hydroxyproline, 3-(C<sub>1-4</sub>alkoxy)proline, 4(C<sub>1-</sub>

4alkoxy)proline, 3,4-dehydroproline, 2,2-dimethyl-4-thiazolidine carboxylic acid, 2,2-dimethyl-4-oxazolidine carboxylic acid, pipercolinic acid, valine, methionine, cysteine, asparagine, serine, threonine,  
5 leucine, tert-leucine, isoleucine, phenylalanine, 1-naphthalanine, 2-naphthalanine, 2-thienylalanine, 3-thienylalanine, [1,2,3,4]-tetrahydroisoquinoline-2-carboxylic acid, aspartic acid-4-C<sub>1-4</sub>alkyl ester and glutamic acid 5-C<sub>1-4</sub>alkyl ester and

10

wherein the second D or L amino acid, is bound to the amino terminus of said first amino acid, and is selected from the group consisting of phenylalanine, 4-benzoylphenylalanine, 4-carboxyphenylalanine, 4-  
15 (carboxyC<sub>1-2</sub>alkyl)phenylalanine, substituted phenylalanine (where the phenyl substituents are independently selected from one or more of C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxy  
20 or C<sub>1-4</sub>alkoxycarbonyl), 3-benzothienylalanine, 4-biphenylalanine, homophenylalanine, octahydroindole-2-carboxylic acid, 2-pyridylalanine, 3-pyridylalanine, 4-thiazolylalanine, 2-thienylalanine, 3-(3-benzothienyl)alanine, 3-thienylalanine, tryptophan,  
25 tyrosine, asparagine, 3-tri-C<sub>1-4</sub>alkylsilylalanine, cyclohexylglycine, diphenylglycine, phenylglycine, methionine sulfoxide, methionine sulfone, 2,2-dicyclohexylalanine, 2-(1-naphthylalanine), 2-(2-naphthylalanine), phenyl substituted phenylalanine  
30 (where the substituents are selected from C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxy or C<sub>1-4</sub>alkoxycarbonyl), aspartic acid, aspartic acid-4-C<sub>1-4</sub>alkyl ester, glutamic acid, glutamic acid-5-C<sub>1-</sub>

4alkyl ester, cycloC<sub>3</sub>-8alkylalanine, substituted  
cycloC<sub>3</sub>-8alkylalanine (where the ring substituents are  
carboxy, C<sub>1</sub>-4 alkyl ester, cycloC<sub>3</sub>-8alkylalanine,  
substituted cycloC<sub>3</sub>-8alkylalanine (where the ring  
5 substituents are carboxy, C<sub>1</sub>-4alkylcarboxy, C<sub>1</sub>-  
4alkoxycarbonyl or aminocarbonyl), 2,2-diphenylalanine  
and all alpha-C<sub>1</sub>-5alkyl of all amino acid derivatives  
thereof, and


10 wherein the amino terminus of said second amino  
acid is unsubstituted or monosubstituted with a member  
of the group consisting of formyl, C<sub>1</sub>-12 alkyl,  
tetrazol-5-yl C<sub>1</sub>-2alkyl, carboxyC<sub>1</sub>-8alkyl,  
carboalkoxyC<sub>1</sub>-4alkyl, phenyl C<sub>1</sub>-4alkyl, substituted  
15 phenylC<sub>1</sub>-4alkyl (where the phenyl substituents or  
independently selected from one or more of, C<sub>1</sub>-4alkyl,  
perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy, hydroxy, halo, amido,  
nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-4dialkylamino, carboxy  
or C<sub>1</sub>-4alkoxycarbonyl), 1,1-diphenylC<sub>1</sub>-4alkyl, C<sub>1</sub>-  
20 6alkoxycarbonyl, phenylC<sub>1</sub>-6alkoxycarbonyl, C<sub>1</sub>-  
2alkylcarbonyl, perfluoro C<sub>1</sub>-4alkylcarbonyl, C<sub>1</sub>-  
4alkylcarbonyl, phenylC<sub>1</sub>-4alkylcarbonyl, substituted  
phenylC<sub>1</sub>-4alkylcarbonyl (where the phenyl substituents  
are independently selected from one or more of C<sub>1</sub>-  
25 4alkyl, perfluoro C<sub>1</sub>-4alkyl, C<sub>1</sub>-4 alkoxy, hydroxy,  
halo, amido, nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-  
4dialkylamino, carboxy or C<sub>1</sub>-4alkoxycarbonyl), 1,1-  
diphenylC<sub>1</sub>-4alkyl, perfluoroC<sub>1</sub>-4alkyl, C<sub>1</sub>-  
4alkoxycarbonyl), 10-camphorsulfonyl, phenylC<sub>1</sub>-  
30 4alkylsulfonyl, substituted phenylC<sub>1</sub>-4alkylsulfonyl, C<sub>1</sub>-  
4alkylsulfinyl, perfluoroC<sub>1</sub>-4alkylsulfinyl,  
phenylsulfinyl, substituted phenylsulfinyl (where the  
phenyl substituents are independently selected from one

or more of, C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxyl or C<sub>1-4</sub>alkoxycarbonyl), phenylC<sub>1-4</sub>alkylsulfinyl, substituted phenylC<sub>1-4</sub>alkylsulfinyl, 1-naphthylsulfonyl, 2-naphthylsulfonyl, substituted naphthylsulfonyl (where the naphthyl substituent is selected from C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxy or C<sub>1-4</sub>alkoxycarbonyl), 1-naphthyl-sulfinyl, 2-naphthylsulfinyl and substituted naphthylsulfinyl (where the naphthyl substituent is selected from C<sub>1-4</sub>alkyl, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, carboxyl or C<sub>1-4</sub>alkoxycarbonyl); R<sub>1</sub> is selected from the group consisting of hydrogen and alkyl; R<sub>2</sub> is selected from the group consisting of amino C<sub>2-5</sub>alkyl, guanidinoC<sub>2-5</sub>alkyl, C<sub>1-4</sub>alkylguanidinoC<sub>2-5</sub>alkyl, diC<sub>1-4</sub>alkylguanidinoC<sub>2-5</sub>alkyl, amidinoC<sub>2-5</sub>alkyl, C<sub>1-4</sub>alkylamidinoC<sub>2-5</sub>alkyl, diC<sub>1-4</sub>alkylamidinoC<sub>2-5</sub>alkyl, C<sub>1-3</sub>alkoxyC<sub>2-5</sub>alkyl, phenyl, substituted phenyl (where the substituents are independently selected from one or more of amino, amidino, guanidino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, halogen, perfluoro C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>1-3</sub>alkoxy or nitro), benzyl, phenyl substituted benzyl (where the substituents are independently selected from one or more of, amino, amidino, guanidino, C<sub>1-4</sub>alkylamino, C<sub>1-4</sub>dialkylamino, halogen, perfluoroC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>1-3</sub>alkoxy or nitro), hydroxyC<sub>2-5</sub>alkyl, C<sub>1-5</sub>alkylaminoC<sub>2-5</sub>alkyl, C<sub>1-5</sub>dialkylaminoC<sub>2-5</sub>alkyl, 4-aminocyclohexylC<sub>0-2</sub>alkyl and C<sub>1-5</sub>alkyl;

p is 0 or 1;

B is

5



10 where n is 0-3, R<sub>3</sub> is H or C<sub>1</sub>-5alkyl and the carbonyl moiety of B is bound to E; E is a heterocycle selected from the group consisting of oxazolin-2-yl, oxazol-2-yl, thiazol-2-yl, thiazol-5-yl, thiazol-4-yl, thiazolin-2-yl, imidazol-2-yl, 4-oxo-2-quinoxalin-2-yl, 15 2-pyridyl, 3-pyridyl, benzo[b]thiophen-2-yl, triazol-4-yl triazol-6-yl, pyrazol-2-yl, 4,5,6,7-tetrahydrobenzothiazol-2-yl, naphtho[2,1-d]thiazol-2-yl, naphtho[1-2-d]thiazol-2-yl quinoxalin-2-yl, isoquinolin-1-yl, isoquinolin-3-yl, benzo [b]furan-2-yl, pyrazin-2-yl, quinazolin-2-yl, isothiazol-5-yl, 20 isothiazol-3-yl, purin-8-yl and a substituted heterocycle where the substituents are selected from C<sub>1</sub>-4, perfluoro C<sub>1</sub>-4alkyl, C<sub>1</sub>-4alkoxy, hydroxy, halo, amido, nitro, amino, C<sub>1</sub>-4alkylamino, C<sub>1</sub>-4dialkylamino, 25 carboxyl, C<sub>1</sub>-4alkoxycarbonyl, hydroxy or phenylC<sub>1</sub>-4 alkylaminocarbonyl, indol-2-yl, benzoxazol-2-yl, benzimidazol-2-yl and benzothiazol-2-yl.

30

What is claimed is:

1. A composition of matter for treating a mammal afflicted with a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a therapeutically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier.
2. A composition of matter for treating a mammal afflicted with a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a therapeutically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier.
3. A composition of matter for preventing in a mammal a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a prophylactically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression, and (b) a pharmaceutically or cosmetically acceptable carrier.
4. A composition of matter for preventing in a mammal a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises (a) a prophylactically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression, and

(b) a pharmaceutically or cosmetically acceptable carrier.

- 5           5.    The composition of claim 1 or 3, wherein the composition comprises an agent which activates the PAR-2 pathway.
- 10           6.    The composition of claim 5, wherein the composition comprises an agent selected from the group consisting of SLIGRL, SAIGRL, SLIGKVD and a serine protease.
- 15           7.    The composition of claim 6, wherein the agent is selected from the group consisting of SLIGRL, trypsin, thrombin and tryptase.
- 20           8.    The composition of claim 2 or 4, wherein the composition comprises an agent which inhibits the PAR-2 pathway.
- 25           9.    The composition of claim 2 or 4, wherein the composition comprises an agent selected from the group consisting of a soybean derivative and a serine protease inhibitor.
- 30           10.   The composition of claim 9, wherein the agent is selected from the group consisting of soybean milk, soybean paste, Compound I, a trypsin inhibitor, a tryptase inhibitor, a thrombin inhibitor and STI.
11.   The composition of claim 1, 2, 3 or 4, wherein the appropriate cells are PAR-2-expressing cells.

12. The composition of claim 11, wherein the appropriate cells are selected from the group consisting of keratinocytes, fibroblasts, and professional phagocytes.
- 5
13. The composition of claim 12, wherein the appropriate cells are keratinocytes.
14. The composition of claim 12, wherein the appropriate cells are fibroblasts.
- 10
15. The composition of claim 12, wherein the appropriate cells are professional phagocytes.
- 15 16. The composition of claim 1, 2, 3 or 4, wherein the disorder is selected from the group consisting of a skin disorder, an immune system disorder, an inflammatory disorder, a respiratory disorder, and a central nervous system disorder.
- 20
17. The composition of claim 16, wherein the disorder is a skin disorder.
18. The composition of claim 16, wherein the disorder is an immune system disorder.
- 25
19. The composition of claim 16, wherein the disorder is an inflammatory disorder.
- 30 20. The composition of claim 16, wherein the disorder is a respiratory disorder.
21. The composition of claim 16, wherein the disorder is a central nervous system disorder.
- 35



22. The composition of claim 1, 2, 3 or 4, wherein the mammal is a human.
23. A method of increasing phagocytosis or ICAM-1 expression in a mammalian cell, comprising contacting the cell with an effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression.
24. A method of decreasing phagocytosis or ICAM-1 expression in a mammalian cell, comprising contacting the cell with an effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.
25. The method of claim 23, wherein the agent activates the PAR-2 pathway.
26. The method of claim 25, wherein the agent is selected from the group consisting of SLIGRL, SAIGRL, SLIGKVD and a serine protease.
27. The method of claim 26, wherein the agent is selected from the group consisting of SLIGRL, trypsin, thrombin and tryptase.
28. The method of claim 24, wherein the agent inhibits the PAR-2 pathway.
29. The method of claim 24, wherein the agent is selected from the group consisting of a soybean derivative and a serine protease inhibitor.
30. The method of claim 29, wherein the agent is selected from the group consisting of soybean

milk, soybean paste, Compound I, a trypsin inhibitor, a tryptase inhibitor, a thrombin inhibitor and STI.

- 5 31. The method of claim 23 or 24, wherein the mammalian cell is a PAR-2-expressing cell.
32. The method of claim 31, wherein the mammalian cell is selected from the group consisting of a  
10 keratinocyte, a fibroblast, and a professional phagocyte.
33. The method of claim 32, wherein the mammalian cell is a keratinocyte.
- 15 34. The method of claim 32, wherein the mammalian cell is a fibroblast.
35. The method of claim 32, wherein the mammalian cell  
20 is a professional phagocyte.
36. The method of claim 23 or 24, wherein the mammalian cell is a human cell.
- 25 37. A method of treating a mammal afflicted with a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a therapeutically effective amount of an agent  
30 that specifically increases phagocytosis or ICAM-1 expression.
38. A method of treating a mammal afflicted with a disorder ameliorated by a decrease in phagocytosis  
35 or ICAM-1 expression in appropriate cells, which

comprises administering to the mammal a therapeutically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.

5

39. A method of preventing in a mammal a disorder ameliorated by an increase in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a prophylactically effective amount of an agent that specifically increases phagocytosis or ICAM-1 expression.

10

40. A method of preventing in a mammal a disorder ameliorated by a decrease in phagocytosis or ICAM-1 expression in appropriate cells, which comprises administering to the mammal a prophylactically effective amount of an agent that specifically decreases phagocytosis or ICAM-1 expression.

15

20

41. The method of claim 37 or 39, wherein the agent activates the PAR-2 pathway.

25

42. The method of claim 41, wherein the agent is selected from the group consisting of SLIGRL, SAIGRL, SLIGKVD and a serine protease.

30

43. The method of claim 42, wherein the agent is selected from the group consisting of SLIGRL, trypsin, thrombin and tryptase.

44. The method of claim 38 or 40, wherein the agent inhibits the PAR-2 pathway.


45. The method of claim 38 or 40, wherein the agent is selected from the group consisting of a soybean derivative and a serine protease inhibitor.
- 5 46. The method of claim 45, wherein the agent is selected from the group consisting of soybean milk, soybean paste, Compound I, a trypsin inhibitor, a tryptase inhibitor, a thrombin inhibitor and STI.
- 10 47. The method of claim 37, 38, 39 or 40, wherein the appropriate cells are PAR-2-expressing cells.
- 15 48. The method of claim 47, wherein the appropriate cells are selected from the group consisting of keratinocytes, fibroblasts, and professional phagocytes.
- 20 49. The method of claim 48, wherein the appropriate cells are keratinocytes.
50. The method of claim 48, wherein the appropriate cells are fibroblasts.
- 25 51. The method of claim 48, wherein the appropriate cells are professional phagocytes.
- 30 52. The method of claim 37, 38, 39 or 40, wherein the disorder is selected from the group consisting of a skin disorder, an immune system disorder, an inflammatory disorder, a respiratory disorder and a central nervous system disorder.
- 35 53. The method of claim 52, wherein the disorder is a skin disorder.

54. The method of claim 52, wherein the disorder is an immune system disorder.
- 5 55. The method of claim 52, wherein the disorder is an inflammatory disorder.
56. The method of claim 52, wherein the disorder is a respiratory disorder.
- 10 57. The method of claim 52, wherein the disorder is a central nervous system disorder.
58. The method of claim 37, 38, 39 or 40, wherein the mammal is a human.
- 15 59. An article of manufacture for administering to a mammal the composition of matter of claim 1, 2, 3 or 4, comprising a solid delivery vehicle having the composition operably affixed thereto.
- 20 60. The article of claim 59, wherein the composition comprises an agent which activates the PAR-2 pathway.
- 25 61. The article of claim 60, wherein the composition comprises an agent selected from the group consisting of SLIGRL, SAIGRL, SLIGKVD and a serine protease.
- 30 62. The article of claim 61, wherein the agent is SLIGRL.

63. The article of claim 59, wherein the composition comprises an agent which inhibits the PAR-2 pathway.
- 5 64. The article of claim 59, wherein the composition comprises an agent selected from the group consisting of a soybean derivative and a serine protease inhibitor.
- 10 65. The article of claim 64, wherein the agent is selected from the group consisting of soybean milk, soybean paste, Compound I, a trypsin inhibitor, a trypsin inhibitor, a thrombin inhibitor and STI.
- 15 66. A method of administering a therapeutic, prophylactic or cosmetic compound to a mammal, comprising administering to the mammal (a) the compound and (b) a composition of matter  
20 comprising a pharmaceutically or cosmetically acceptable carrier and an agent that specifically increases phagocytosis in an amount sufficient to increase phagocytosis in cells where uptake of the compound is desired, wherein the composition is  
25 administered prior to and/or concurrently with the administration of the compound.
67. The method of claim 66, wherein the composition comprises an agent which activates the PAR-2  
30 pathway.
68. The method of claim 67, wherein the composition comprises an agent selected from the group consisting of SLIGRL, SAIGRL, SLIGKVD and a serine  
35 protease.


69. The method of claim 68, wh rein the agent is  
SLIGRL.

**FIG. 1**







**FIG. 2**




**FIG. 3**




**FIG. 4A**




**FIG. 4B**






**FIG. 5A**


**FIG. 5B**






**FIG. 6A**

# FIG. 6B






**FIG. 7A**



**FIG. 7B**




Control

SLIGRL

**FIG. 7C**

Control




STI

**FIG. 8**


Control




SLIGRL




SH00230






**FIG. 9**

**FIG. 10**



**FIG. 11**



# INTERNATIONAL SEARCH REPORT

Int. l. Application No  
PCT/US 98/26255

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6    A61K38/08    A61K38/48    A61K38/55    A61K38/17    A61K38/05 //A61K7/00, A61K9/127				
According to International Patent Classification (IPC) or to both national classification and IPC				
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6    A61K				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)				
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>				
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	EP 0 421 021 A (MUCOS EMULSIONSGESELLSCHAFT M.B.H.) 10 April 1991 see page 2, line 1 - line 4; claims 1,10 see page 1, line 15 - line 23 ---	1, 3, 23, 37, 39, 58, 66		
X	WO 96 24371 A (PHAIRSON MEDICAL, INC.) 15 August 1996  see page 6, line 16 - page 9, line 25; claims see page 13, line 21 - line 28 --- -/--	2, 4, 16-20, 24, 38, 40, 52-56, 58		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;"><input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.</td> <td style="width: 50%; border: none;"><input checked="" type="checkbox"/> Patent family members are listed in annex.</td> </tr> </table>			<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.	<input checked="" type="checkbox"/> Patent family members are listed in annex.
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.	<input checked="" type="checkbox"/> Patent family members are listed in annex.			
* Special categories of cited documents :				
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;">                     *A* document defining the general state of the art which is not considered to be of particular relevance                      *E* earlier document but published on or after the international filing date                      *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)                      *O* document referring to an oral disclosure, use, exhibition or other means                      *P* document published prior to the international filing date but later than the priority date claimed                 </td> <td style="width: 50%; border: none;">                     *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention                      *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone                      *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.                      *S* document member of the same patent family                 </td> </tr> </table>			*A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *S* document member of the same patent family
*A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *S* document member of the same patent family			
Date of the actual completion of the international search  <p style="text-align: center; font-weight: bold;">22 April 1999</p>	Date of mailing of the international search report  <p style="text-align: center; font-weight: bold;">06/05/1999</p>			
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016	Authorized officer  <p style="text-align: center; font-weight: bold;">Ryckebosch, A</p>			



INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/26255

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>C.K. DERIAN ET L.: "DIFFERENTIAL REGULATION OF HUMAN KERATINOCYTE GROWTH AND DIFFERENTIATION BY A NOVEL FAMILY OF PROTEASE-ACTIVATED RECEPTORS." CELL GROWTH &amp; DIFFERENTIATION, vol. 8, no. 7, July 1997, pages 743-749, XP002100833 BALTIMORE, MD, US cited in the application see page 743, abstract ---</p>	1-69
A	<p>R.J. SANTULLI ET AL.: "EVIDENCE FOR THE PRESENCE OF A PROTEASE-ACTIVATED RECEPTOR DISTINCT FROM THE THROMBIN RECEPTOR IN HUMAN KERATINOCYTES." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 92, September 1995, pages 9151-9155, XP002100834 WASHINGTON US see page 9151, abstract ---</p>	1-69
A	<p>M. MOLINO ET AL.: "INTERACTIONS OF MAST CELL TRYPTASE WITH THROMBIN RECEPTORS AND PAR-2" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 272, no. 7, 14 February 1997, pages 4043-4049, XP002094579 BALTIMORE, MD, US see page 4043, abstract -----</p>	1-69

INTERNATIONAL SEARCH REPORT

...ernational application No.

PCT/US 98/26255

**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: 23-36, 37-58, 66-69  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 23-36 (all as far as being related to an in vivo method), 37-58 and 66-69 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2.  Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3.  Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.  As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.  No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Jonal Application No

PCT/US 98/26255

Patent document cited in search report	A	Publication date	Patent family member(s)	Publication date
EP 421021	A	10-04-1991	NONE	
WO 9624371	A	15-08-1996	AU 4917096 A BR 9607506 A CA 2212533 A CN 1181018 A CZ 9702541 A EP 0810875 A JP 11502102 T NO 973627 A PL 324085 A ZA 9601030 A	27-08-1996 23-12-1997 15-08-1996 06-05-1998 18-03-1998 10-12-1997 23-02-1999 07-10-1997 11-05-1998 29-08-1996